

Measures in ternary homogeneous structures

By Paolo Marimon, p.marimon19@imperial.ac.uk

① Measures on countable graphs and hypergraphs

Measures are important in model theory since

- they give a good notion of size on infinite structures;
- they often capture how a structure can be obtained by random constructions (e.g. the **random graph**);
- they naturally arise in infinite structures approximating the behaviour of a class of finite structures (e.g. **pseudofinite fields**).

We focus on probability measures arising in highly symmetrical countable **graphs** and **hypergraphs**.

The 3-hypergraphs that we study exhibit oddly-behaved measures which serve as counterexamples to many conjectures in the field.

② Definable sets in graphs and hypergraphs

We focus on **definable sets** in graphs and hypergraphs. For example, for a graph G and $a, b \in G$,

$$E(x, a) \wedge E(x, b)$$

defines the set of vertices $x \in G$ joined to both a and b . A **3-hypergraph** has a 3-hyperedge relation $R(x, y, z)$.

We work with **homogeneous** structures: any isomorphism between finite substructures extends to an automorphism of the whole structure.

③ Invariant Keisler measures

A **Keisler measure** on a structure \mathcal{M} is a finitely additive **probability** measure on its definable subsets. We are interested in measures **invariant under automorphisms**:

$$\mu(X) = \mu(\sigma(X)) \text{ for } \sigma \in \text{Aut}(M).$$

For $d \in M, A \subset M, \mu(d/A)$ is the measure of the intersection of all sets containing d defined with parameters from A .

④ Example: measures in homogeneous graphs

Countable homogeneous graphs have been classified in [5]. Two important examples:

- the **random graph**. The unique countable homogeneous graph containing every finite graph as a subgraph;
- the **generic triangle-free graph**. The unique countable homogeneous graph containing every finite triangle-free graph.

In a countable homogeneous graph, the measure $\mu(d/A)$ **does not depend on the structure of A**

For example, for $a, b \in G$ forming an edge and $a', b' \in G$ not forming an edge,

$$\mu(E(x, a) \wedge E(x, b)) = \mu(E(x, a') \wedge E(x, b')).$$



This is due to a very general and powerful principle that holds in this context [4]:

Stationarity: Given a and b independent,

$$\mu(\phi(x, a) \wedge \psi(x, b))$$

only depends on the structure of a and b respectively, and not on the structure of ab .

Stationarity and some ergodic theory allow for a full classification of invariant Keisler measures on homogeneous countable graphs [1].

⑤ Higher stationarity?

Given a, b, c satisfying adequate independence assumptions, can we prove that


$$\mu(\phi(x, ab) \wedge \psi(x, ac) \wedge \chi(x, bc))$$

only depends on the structure of the pairs ab, ac, bc and not on the structure of the whole triplet?

A version of this **higher stationarity** holds in many structures (e.g. homogeneous graphs). Moreover it was suggested by hypergraph regularity results in pseudofinite fields [2].

⑥ The universal homogeneous two-graph has a unique measure

A **two-graph** is a 3-hypergraph such that every four vertices have an even number of hyperedges.



The **universal homogeneous two-graph** \mathcal{G} is the unique countable 3-hypergraph containing every finite two-graph as a substructure.

A unique measure

There is a unique invariant Keisler measure for \mathcal{G} . Given d, a_1, \dots, a_n distinct, we have that

$$\mu(d/a_1, \dots, a_n) = \left(\frac{1}{2}\right)^{n-1}.$$

The universal homogeneous two-graph gives a **counterexample to higher stationarity**:

Take $a, b, c \in \mathcal{G}$ not forming a hyperedge. Then,


$$\mu(R(x, ab) \wedge R(x, ac) \wedge R(x, bc)) = 0,$$

because if there was any such x , there would be four vertices with three hyperedges. Meanwhile, for $a', b', c' \in \mathcal{G}$ forming a hyperedge,

$$\mu(R(x, a'b') \wedge R(x, a'c') \wedge R(x, b'c')) = \frac{1}{4}.$$

⑦ The generic tetrahedron-free 3-hypergraph is not MS-measurable

A **tetrahedron** is a 3-hypergraph of four vertices such that each three of them form a hyperedge.




The **generic tetrahedron-free 3-hypergraph** \mathcal{H} is the unique countable homogeneous 3-hypergraph containing every finite tetrahedron-free 3-hypergraph as a substructure.

An MS-measurable structure [6] has a well-behaved system of a dimension and measures. It was believed that various properties satisfied by \mathcal{H} jointly implied MS-measurability [3]. However,

No "well-behaved" system of measures

The generic tetrahedron-free 3-hypergraph is not MS-measurable.

The trick behind the proof is **forcing higher stationarity**: fix d and consider a, b, c such that each pair forms a hyperedge with it. abc cannot form a hyperedge since we **avoid tetrahedrons!**



Hence, for any formulas, and μ $\text{Aut}(H/d)$ -invariant:

$$\mu(\phi(x, ab, d) \wedge \psi(x, ac, d) \wedge \chi(x, bc, d))$$

is entirely determined by the structure of each triplet containing d and two vertices from abc .

⑧ References

- [1] M. H. Albert. "Measures on the random graph". In: *J London Math Soc* 50.3 (1994), pp. 417-429.
- [2] A. Chevalier and E. Levi. *An Algebraic Hypergraph Regularity Lemma*. 2022. arXiv: 2204.01158 [math.CO].
- [3] R. Elwes and D. Macpherson. "A survey of asymptotic classes and measurable structures". In: *Model Theory with Applications to Algebra and Analysis*. Vol. 2. London Mathematical Society Lecture Note Series. CUP, 2008, pp. 125-160.
- [4] E. Hrushovski. "Stable Group Theory and Approximate subgroups". In: *J Am Math Soc* 25.1 (2012), pp. 189-243.
- [5] A. H. Lachlan and R. E. Woodrow. "Countable Ultrahomogeneous Undirected Graphs". In: *Trans Am Math Soc* 262.1 (1980), pp. 51-94.
- [6] D. Macpherson and C. Steinhorn. "One-dimensional asymptotic classes of finite structures". In: *Trans Am Math Soc* 360.1 (2008), pp. 411-448.