

## References

- Beste, D.J., *et al.* The genetic requirements for fast and slow growth in mycobacteria. *PLoS One* **4**, e5349 (2009).
- Cantone, I., *et al.* A yeast synthetic network for in vivo assessment of reverse-engineering and modeling approaches. *Cell* **137**, 172-181 (2009).
- Caron, E. & Hall, A. Identification of two distinct mechanisms of phagocytosis controlled by different Rho GTPases. *Science* **282**, 1717-1721 (1998).
- Ehrt, S., *et al.* Reprogramming of the macrophage transcriptome in response to interferon-gamma and Mycobacterium tuberculosis: signaling roles of nitric oxide synthase-2 and phagocyte oxidase. *J Exp Med* **194**, 1123-1140 (2001).
- Grode, L., *et al.* Increased vaccine efficacy against tuberculosis of recombinant *Mycobacterium bovis* bacille Calmette-Guérin mutants that secrete listeriolysin. *J Clin Invest.* **115**, 2472-9 (2005)
- Groves, E., *et al.* Sequestering of Rac by the Yersinia effector YopO blocks Fcγ receptor-mediated phagocytosis. *J Biol Chem* **285**, 4087-4098 (2010).
- Khorashad, J.S., *et al.* The level of BCR-ABL1 kinase activity before treatment does not identify chronic myeloid leukemia patients who fail to achieve a complete cytogenetic response on imatinib. *Haematologica* **94**, 861-864 (2009).
- Kuijl, C., *et al.* Intracellular bacterial growth is controlled by a kinase network around PKB/AKT1. *Nature* **450**, 725-730 (2007).
- Ly, K.T. & Casanova, J.E. Abelson tyrosine kinase facilitates Salmonella enterica serovar Typhimurium entry into epithelial cells. *Infect Immun* **77**, 60-69 (2009).
- Mochizuki, Y. & Majerus, P.W. Characterization of myotubularin-related protein 7 and its binding partner, myotubularin-related protein 9. *Proc Natl Acad Sci U S A* **100**, 9768-9773 (2003).
- Paul, S., Laochumroonvorapong, P. & Kaplan, G. Comparable growth of virulent and avirulent *Mycobacterium tuberculosis* in human macrophages in vitro. *J Infect Dis* **174**, 105-112 (1996).
- Shin HW, *et al.* An enzymatic cascade of Rab5 effectors regulates phosphoinositide turnover in the endocytic pathway. *J Cell Biol* **170**: 607-618 (2005).
- Slavkov, I., Gjorgjoski, V., Struyf, J. & Dzeroski, S. Finding explained groups of time-course gene expression profiles with predictive clustering trees. *Mol Biosyst* **6**, 729-740 (2010).
- Stewart, G.R., Patel, J., Robertson, B.D., Rae, A. & Young, D.B. Mycobacterial mutants with defective control of phagosomal acidification. *PLoS Pathog* **1**, 269-278 (2005).
- Tashkova et al. (2010) Parameter estimation in an endocytosis model with bio-inspired optimization algorithms. In Filipic, B. & Silc, J., eds, *Proceedings of the Fourth International Conference on Bioinspired Optimization Methods and their Applications* (BIOMA 2010), p67-82. Jožef Stefan Institute, Ljubljana.
- Todorovski & Dzeroski (2009) Inductive process-based modeling of endocytosis from time-series data. In: DŽEROSKI, Sašo (ed.), GEURTS, Pierre (ed.), ROUSU, Juho (ed.). *Machine Learning in Systems Biology : Proceedings of the Third International Workshop, September 5-6, 2009, Ljubljana, Slovenia*, (Julkaisusarja - Helsingin yliopisto. Tietojenkäsittelytieteen laitos, report B-2009-1). Helsinki University, Department of Computer Science, 2009, p175-176.
- Van Etten, R.A., *et al.* The COOH terminus of the c-Abl tyrosine kinase contains distinct F- and G-actin binding domains with bundling activity. *J Cell Biol* **124**, 325-340 (1994).
- Verreck, F.A., *et al.* Human IL-23-producing type 1 macrophages promote but IL-10-producing type 2 macrophages subvert immunity to (myco)bacteria. *Proc Natl Acad Sci U S A* **101**, 4560-4565 (2004).
- Verreck, F.A., de Boer, T., Langenberg, D.M., van der Zanden, L. & Ottenhoff, T.H. Phenotypic and functional profiling of human proinflammatory type-1 and anti-inflammatory type-2 macrophages in response to microbial antigens and IFN-gamma- and CD40L-mediated costimulation. *J Leukoc Biol* **79**, 285-293 (2006).
- Wang, Y., Miller, A.L., Mooseker, M.S. & Koleske, A.J. The Abl-related gene (Arg) nonreceptor tyrosine kinase uses two F-actin-binding domains to bundle F-actin. *Proc Natl Acad Sci U S A* **98**, 14865-14870 (2001).
- Wiedemann, A., *et al.* Two distinct cytoplasmic regions of the beta2 integrin chain regulate RhoA function during phagocytosis. *J Cell Biol* **172**, 1069-1079 (2006).
- Yeung T & Grinstein S. Lipid signaling and the modulation of surface charge during phagocytosis. *Immunol Rev* **219**: 17-36 (2007).